

Vectors: The DOT

What is *the DOT Product*

The dot product is a way of 'multiplying' two vectors together. The result is not a vector but a number. This number is usually referred to as *the DOT product* of the two vectors. Moreover, by itself and at first glance, the dot product appears to be a clumsy nobody. However, spend some time with the DOT and you will find it gives a clean and graceful way to the obtain magnitude of a vector, 'the distance' between two vectors, the angle between two vectors, a simple test to determine if two vectors are perpendicular, and elegant way to project a vector onto another, AND grand generalization to spaces way beyond \mathbf{R}^2 , way beyond. Having said that, the next task at hand is to actually define the dot product. Let us start with vectors in \mathbf{R}^2 . For any vector $\mathbf{v} = \langle a, b \rangle$ and any other vector $\mathbf{u} = \langle c, d \rangle$, then the dot product is written as $\mathbf{v} \cdot \mathbf{u}$ and is defined as follows:

$$\langle a, b \rangle \cdot \langle c, d \rangle = ac + bd$$

some famous *DOT product properties*


The properties below are precisely what makes the dot product special. It is these properties exactly that allow us to find magnitudes, distances, angles between vectors, all all the other perks that come with being a dot product. This is important to note because, on a different occasion, we may meet a different 'dot product', but so long as it meets these conditions it too will yield a magnitude for vectors, distance, angles, projections, etc... Said differently, anyone who wants to be a dot product has to meet these conditions, and in doing so, will bring with it all the great perks that come with being a dot product.⁰

$\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$	<i>commutes</i>
$(\mathbf{u} + \mathbf{w}) \cdot \mathbf{v} = \mathbf{u} \cdot \mathbf{v} + \mathbf{w} \cdot \mathbf{v}$	<i>distributes</i>
$\mathbf{u} \cdot (c\mathbf{v}) = c(\mathbf{u} \cdot \mathbf{v})$	<i>pull constant</i>
$\mathbf{u} \cdot \mathbf{u} \geq 0$	<i>self dot pos</i>
$\mathbf{u} \cdot \mathbf{u} = 0 \iff \mathbf{u} = \mathbf{0}$	<i>self dot 0</i>

Magnitude by *the DOT*


Observe what happens when a vector is dotted with itself.

$$\mathbf{v} \cdot \mathbf{v} = \langle a, b \rangle \cdot \langle a, b \rangle = a^2 + b^2$$



'Distance' by *the DOT*

It should be emphasized that the vectors we are working with are defined by their direction and magnitude, NOT by their position. Even without a fixed position we can still define some sort of *distance* between two vectors. This is important because further study of mathematics will lead to great generalizations of the *distance* concept. For the moment, we will be content to note we can measure a *distance* between two vectors in the following way:



1. Vector Dot Product

(a) Compute the following DOT product:

$$\langle 1, 2 \rangle \cdot \langle 3, 5 \rangle$$

Solution:

$$\begin{aligned}\langle 1, 2 \rangle \cdot \langle 3, 5 \rangle &= (1)(3) + (2)(5) && \text{(def of the DOT)} \\ &= 3 + 10 && \text{(by Calc)} \\ &= 13 && \text{(by Calc)}\end{aligned}$$

(b) Compute the following DOT product:

$$\langle 3, 5 \rangle \cdot \langle 3, 5 \rangle$$

Solution:

$$\begin{aligned}\langle 3, 5 \rangle \cdot \langle 3, 5 \rangle &= (3)(3) + (5)(5) && \text{(def of the DOT)} \\ &= 9 + 25 && \text{(by Calc)} \\ &= 34 && \text{(by Calc)}\end{aligned}$$

(c) Compute the following DOT product:

$$\langle -3, 2 \rangle \cdot \langle 1, 2 \rangle$$

Solution:

$$\begin{aligned}\langle -3, 2 \rangle \cdot \langle 1, 2 \rangle &= (-3)(1) + (2)(2) && \text{(def of the DOT)} \\ &= -3 + 4 && \text{(by Calc)} \\ &= 1 && \text{(by Calc)}\end{aligned}$$

(d) Compute the following DOT product:

$$\langle 1, 4 \rangle \cdot \langle -3, -5 \rangle$$

Solution:

$$\begin{aligned}\langle 1, 4 \rangle \cdot \langle -3, -5 \rangle &= (1)(-3) + (4)(-5) && \text{(def of the DOT)} \\ &= -3 + -20 && \text{(by Calc)} \\ &= -23 && \text{(by Calc)}\end{aligned}$$

(e) Compute the following DOT product:

$$\langle 2, 3 \rangle \cdot \langle 3, -3 \rangle$$

Solution:

$$\begin{aligned}\langle 2, 3 \rangle \cdot \langle 3, -3 \rangle &= (2)(3) + (3)(-3) && \text{(def of the DOT)} \\ &= 6 + -9 && \text{(by Calc)} \\ &= -3 && \text{(by Calc)}\end{aligned}$$

(f) Compute the following DOT product:

$$\langle 1, -4 \rangle \cdot \langle 3, 7 \rangle$$

Solution:

$$\begin{aligned}\langle 1, -4 \rangle \cdot \langle 3, 7 \rangle &= (1)(3) + (-4)(7) && \text{(def of the DOT)} \\ &= 3 + -28 && \text{(by Calc)} \\ &= -25 && \text{(by Calc)}\end{aligned}$$

(g) $\langle 2, 1, 3 \rangle \cdot \langle 1, 2, 1 \rangle$

Solution: $2 + 2 + 3 = 7$

2. Vector Dot Product

(a) Compute the following DOT product:

$$(\mathbf{i} + 2\mathbf{j}) \cdot (\mathbf{0i} + 5\mathbf{j})$$

Solution:

$$\begin{aligned}(\mathbf{i} + 2\mathbf{j}) \cdot (\mathbf{0i} + 5\mathbf{j}) &= (1 \langle 1, 0 \rangle + 2 \langle 0, 1 \rangle) \cdot (0 \langle 1, 0 \rangle + 5 \langle 0, 1 \rangle) && \text{(def of } \mathbf{i} \text{ and } \mathbf{j}) \\ &= \langle 1, 2 \rangle \cdot \langle 0, 5 \rangle && \text{(by Inspection)} \\ &= (1)(0) + (2)(5) && \text{(def of the DOT)} \\ &= 0 + 10 && \text{(by Calc)} \\ &= 10 && \text{(by Calc)}\end{aligned}$$

(b) Compute the following DOT product:

$$(3\mathbf{i} + 0\mathbf{j}) \cdot (3\mathbf{i} + 5\mathbf{j})$$

Solution:

$$\begin{aligned}
 (3\mathbf{i} + 0\mathbf{j}) \cdot (3\mathbf{i} + 5\mathbf{j}) &= (3 \langle 1, 0 \rangle + 0 \langle 0, 1 \rangle) \cdot (3 \langle 1, 0 \rangle + 5 \langle 0, 1 \rangle) && \text{(def of } \mathbf{i} \text{ and } \mathbf{j}) \\
 &= \langle 3, 0 \rangle \cdot \langle 3, 5 \rangle && \text{(by Inspection)} \\
 &= (3)(3) + (0)(5) && \text{(def of } \textit{the DOT}) \\
 &= 9 + 0 && \text{(by Calc)} \\
 &= 9 && \text{(by Calc)}
 \end{aligned}$$

(c) Compute the following DOT product:

$$(-3\mathbf{i} + 2\mathbf{j}) \cdot (\mathbf{i} + 2\mathbf{j})$$

Solution:

$$\begin{aligned}
 (-3\mathbf{i} + 2\mathbf{j}) \cdot (\mathbf{i} + 2\mathbf{j}) &= (-3 \langle 1, 0 \rangle + 2 \langle 0, 1 \rangle) \cdot (1 \langle 1, 0 \rangle + 2 \langle 0, 1 \rangle) && \text{(def of } \mathbf{i} \text{ and } \mathbf{j}) \\
 &= \langle -3, 2 \rangle \cdot \langle 1, 2 \rangle && \text{(by Inspection)} \\
 &= (-3)(1) + (2)(2) && \text{(def of } \textit{the DOT}) \\
 &= -3 + 4 && \text{(by Calc)} \\
 &= 1 && \text{(by Calc)}
 \end{aligned}$$

(d) Compute the following DOT product:

$$(\mathbf{i} + 4\mathbf{j}) \cdot (-3\mathbf{i} + -5\mathbf{j})$$

Solution:

$$\begin{aligned}
 (\mathbf{i} + 4\mathbf{j}) \cdot (-3\mathbf{i} + -5\mathbf{j}) &= (1 \langle 1, 0 \rangle + 4 \langle 0, 1 \rangle) \cdot (-3 \langle 1, 0 \rangle + -5 \langle 0, 1 \rangle) && \text{(def of } \mathbf{i} \text{ and } \mathbf{j}) \\
 &= \langle 1, 4 \rangle \cdot \langle -3, -5 \rangle && \text{(by Inspection)} \\
 &= (1)(-3) + (4)(-5) && \text{(def of } \textit{the DOT}) \\
 &= -3 + -20 && \text{(by Calc)} \\
 &= -23 && \text{(by Calc)}
 \end{aligned}$$

3. Vector Dot Product to find magnitude

(a) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 1, 2 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 1, 2 \rangle \cdot \langle 1, 2 \rangle && \text{(given)} \\
 &= (1)(1) + (2)(2) && \text{(def of the DOT)} \\
 &= 1 + 4 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 5 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{5} \\
 & \|\vec{v}\| \approx 2.24
 \end{aligned}$$

(b) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 3, 5 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 3, 5 \rangle \cdot \langle 3, 5 \rangle && \text{(given)} \\
 &= (3)(3) + (5)(5) && \text{(def of the DOT)} \\
 &= 9 + 25 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 34 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{34} \\
 & \|\vec{v}\| \approx 5.83
 \end{aligned}$$

(c) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle -3, 2 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle -3, 2 \rangle \cdot \langle -3, 2 \rangle && \text{(given)} \\
 &= (-3)(-3) + (2)(2) && \text{(def of the DOT)} \\
 &= 9 + 4 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 13 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{13} \\
 & \|\vec{v}\| \approx 3.61
 \end{aligned}$$

(d) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 1, 4 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 1, 4 \rangle \cdot \langle 1, 4 \rangle && \text{(given)} \\
 &= (1)(1) + (4)(4) && \text{(def of the DOT)} \\
 &= 1 + 16 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 17 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{17} \\
 & \|\vec{v}\| \approx 4.12
 \end{aligned}$$

(e) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 2, 3 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 2, 3 \rangle \cdot \langle 2, 3 \rangle && \text{(given)} \\
 &= (2)(2) + (3)(3) && \text{(def of the DOT)} \\
 &= 4 + 9 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 13 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{13} \\
 & \|\vec{v}\| \approx 3.61
 \end{aligned}$$

(f) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 1, -4 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 1, -4 \rangle \cdot \langle 1, -4 \rangle && \text{(given)} \\
 &= (1)(1) + (-4)(-4) && \text{(def of the DOT)} \\
 &= 1 + 16 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 17 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{17} \\
 & \|\vec{v}\| \approx 4.12
 \end{aligned}$$

(g) Use the dot product to find the magnitude of the following vector:

$$\vec{v} = \langle 3, -4 \rangle$$

Solution:

$$\begin{aligned}
 \|\vec{v}\|^2 &= \vec{v} \cdot \vec{v} && \text{(famous DOT product property)} \\
 &= \langle 3, -4 \rangle \cdot \langle 3, -4 \rangle && \text{(given)} \\
 &= (3)(3) + (-4)(-4) && \text{(def of the DOT)} \\
 &= 9 + 16 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v}\|^2 = 25 && \text{(by Calc)} \\
 \dots\text{then.. (assuming norm is positive)}\dots & \|\vec{v}\| = \sqrt{25} \\
 & \|\vec{v}\| \approx 5
 \end{aligned}$$

4. **Vector Dot Product to find 'distance'**

The DOT product provides a way to define the 'distance' between two vectors, \vec{v} and \vec{w} . So long as we can define subtraction of the vectors we can define the distance between them as follows:

$$dist(\vec{v}, \vec{w}) = \|\vec{v} - \vec{w}\| = \sqrt{(\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})}$$

(a) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 1, 2 \rangle \quad \vec{v} = \langle 3, 5 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle 2, 3 \rangle$

$$\begin{aligned}
 \|\vec{v} - \vec{w}\|^2 &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) && \text{(famous DOT product property)} \\
 &= \langle 2, 3 \rangle \cdot \langle 2, 3 \rangle && \text{(from given)} \\
 &= (2)(2) + (3)(3) && \text{(def of the DOT)} \\
 &= 4 + 9 && \text{(by Calc)} \\
 \dots\text{thus}\dots & \|\vec{v} - \vec{w}\|^2 = 13 && \text{(by Calc)} \\
 \dots\text{then.. (assuming the distance is positive)}\dots & \|\vec{v} - \vec{w}\| = \sqrt{13} \\
 & \|\vec{v} - \vec{w}\| \approx 3.61
 \end{aligned}$$

(b) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle -3, 2 \rangle \quad \vec{v} = \langle 1, 2 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle 4, 0 \rangle$

$$\|\vec{v} - \vec{w}\|^2 = (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})$$

(famous DOT product property)

$$= \langle 4, 0 \rangle \cdot \langle 4, 0 \rangle \quad (\text{from given})$$

$$= (4)(4) + (0)(0) \quad (\text{def of the DOT})$$

$$= 16 + 0 \quad (\text{by Calc})$$

$$\dots\text{thus}\dots \quad \|\vec{v} - \vec{w}\|^2 = 16 \quad (\text{by Calc})$$

$$\dots\text{then.. (assuming the distance is positive)}\dots \quad \|\vec{v} - \vec{w}\| = \sqrt{16}$$

$$\|\vec{v} - \vec{w}\| \approx 4$$

(c) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 1, 4 \rangle \quad \vec{v} = \langle -3, -5 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle -4, -9 \rangle$

$$\|\vec{v} - \vec{w}\|^2 = (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})$$

(famous DOT product property)

$$= \langle -4, -9 \rangle \cdot \langle -4, -9 \rangle \quad (\text{from given})$$

$$= (-4)(-4) + (-9)(-9) \quad (\text{def of the DOT})$$

$$= 16 + 81 \quad (\text{by Calc})$$

$$\dots\text{thus}\dots \quad \|\vec{v} - \vec{w}\|^2 = 97 \quad (\text{by Calc})$$

$$\dots\text{then.. (assuming the distance is positive)}\dots \quad \|\vec{v} - \vec{w}\| = \sqrt{97}$$

$$\|\vec{v} - \vec{w}\| \approx 9.85$$

(d) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 2, 3 \rangle \quad \vec{v} = \langle 3, -3 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle 1, -6 \rangle$

$$\|\vec{v} - \vec{w}\|^2 = (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w})$$

(famous DOT product property)

$$= \langle 1, -6 \rangle \cdot \langle 1, -6 \rangle \quad (\text{from given})$$

$$= (1)(1) + (-6)(-6) \quad (\text{def of the DOT})$$

$$= 1 + 36 \quad (\text{by Calc})$$

$$\dots\text{thus}\dots \quad \|\vec{v} - \vec{w}\|^2 = 37 \quad (\text{by Calc})$$

$$\dots\text{then.. (assuming the distance is positive)}\dots \quad \|\vec{v} - \vec{w}\| = \sqrt{37}$$

$$\|\vec{v} - \vec{w}\| \approx 6.08$$

(e) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 1, -4 \rangle \quad \vec{v} = \langle 3, 7 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle 2, 11 \rangle$

$$\begin{aligned} \|\vec{v} - \vec{w}\|^2 &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) && \text{(famous DOT product property)} \\ &= \langle 2, 11 \rangle \cdot \langle 2, 11 \rangle && \text{(from given)} \\ &= (2)(2) + (11)(11) && \text{(def of the DOT)} \\ &= 4 + 121 && \text{(by Calc)} \end{aligned}$$

$$\dots\text{thus}\dots \quad \|\vec{v} - \vec{w}\|^2 = 125 \quad \text{(by Calc)}$$

$$\begin{aligned} \dots\text{then}\dots \text{ (assuming the distance is positive)}\dots \quad \|\vec{v} - \vec{w}\| &= \sqrt{125} \\ \|\vec{v} - \vec{w}\| &\approx 11.18 \end{aligned}$$

(f) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 3, -4 \rangle \quad \vec{v} = \langle 3, 7 \rangle$$

Solution: First note that $\vec{v} - \vec{w} = \langle 0, 11 \rangle$

$$\begin{aligned} \|\vec{v} - \vec{w}\|^2 &= (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) && \text{(famous DOT product property)} \\ &= \langle 0, 11 \rangle \cdot \langle 0, 11 \rangle && \text{(from given)} \\ &= (0)(0) + (11)(11) && \text{(def of the DOT)} \\ &= 0 + 121 && \text{(by Calc)} \end{aligned}$$

$$\dots\text{thus}\dots \quad \|\vec{v} - \vec{w}\|^2 = 121 \quad \text{(by Calc)}$$

$$\begin{aligned} \dots\text{then}\dots \text{ (assuming the distance is positive)}\dots \quad \|\vec{v} - \vec{w}\| &= \sqrt{121} \\ \|\vec{v} - \vec{w}\| &\approx 11 \end{aligned}$$

(g) Use the dot product to find the 'distance' between the indicated vectors.

$$\vec{w} = \langle 1, -3, 5, 0, 6 \rangle \quad \vec{v} = \langle 5, 2, -10, 3, 1 \rangle$$

Solution: don't be afraid, don't google it, don't ask anyone.. just you and the problem.. its on..., if it knocks you down, just get up... don't let it beat you..

5. **Vector Dot Product to find 'angle' between two vectors**

(a) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 1, 2 \rangle \quad \vec{v} = \langle 3, -5 \rangle$$

Solution: Let us assume the angle between the vectors is between 0 and 180° Suppose we call such angle 'θ', then....


$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 1, 2 \rangle \cdot \langle 3, -5 \rangle}{\|\langle 1, 2 \rangle\| \|\langle 3, -5 \rangle\|} \quad (\text{given}) \\ &\approx \frac{-7}{(2.236)(5.831)} \approx \frac{-7}{13.038} \quad (\text{by Calc}) \end{aligned}$$

....thus.... $\cos \theta = -0.54$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$).... $\theta \approx \cos^{-1}(-0.54)$

$$\theta \approx 2.14 \text{ radians}$$

$$\approx 122.6^\circ$$



(b) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle -3, 2 \rangle \quad \vec{v} = \langle 3, 5 \rangle$$

Solution: Let us assume the angle between the vectors is between 0 and 180° Suppose we call such angle

' θ ', then....

$$\cos \theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property})$$


$$= \frac{\langle -3, 2 \rangle \cdot \langle 3, 5 \rangle}{\|\langle -3, 2 \rangle\| \|\langle 3, 5 \rangle\|} \quad (\text{given})$$

$$\approx \frac{1}{(3.606)(5.831)} \approx \frac{1}{21.027} \quad (\text{by Calc})$$

....thus.... $\cos \theta = 0.05$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$).... $\theta \approx \cos^{-1}(0.05)$

$$\theta \approx 1.52 \text{ radians}$$

$$\approx 87.1^\circ$$


(c) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 1, 4 \rangle \quad \vec{v} = \langle -3, -5 \rangle$$

Solution: Let us assume the angle between the vectors is between 0 and 180° Suppose we call such angle ' θ ', then....

$$\cos \theta = \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property})$$

$$= \frac{\langle 1, 4 \rangle \cdot \langle -3, -5 \rangle}{\|\langle 1, 4 \rangle\| \|\langle -3, -5 \rangle\|} \quad (\text{given})$$


$$\approx \frac{-23}{(4.123)(5.831)} \approx \frac{-23}{24.041} \quad (\text{by Calc})$$

....thus.... $\cos \theta = -0.96$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$).... $\theta \approx \cos^{-1}(-0.96)$

$$\theta \approx 2.86 \text{ radians}$$

$$\approx 163.9^\circ$$



(d) Use the dot product to find the 'angle' between the indicated vectors.


$$\vec{w} = \langle 2, 3 \rangle \quad \vec{v} = \langle 3, -3 \rangle$$

Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 2, 3 \rangle \cdot \langle 3, -3 \rangle}{\|\langle 2, 3 \rangle\| \|\langle 3, -3 \rangle\|} \quad (\text{given}) \\ &\approx \frac{-3}{(3.606)(4.243)} \approx \frac{-3}{15.3} \quad (\text{by Calc}) \end{aligned}$$

$$\dots \text{thus} \dots \quad \cos \theta = -0.2 \quad (\text{by Calc})$$

$$\begin{aligned} \dots \text{then} \dots (\text{assume the sought angle is } 0 \leq \theta \leq 180^\circ) \dots & \quad \theta \approx \cos^{-1}(-0.2) \\ & \quad \theta \approx 1.77 \text{ radians} \\ & \quad \approx 101.4^\circ \end{aligned}$$



(e) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 3, 7 \rangle \quad \vec{v} = \langle 1, -4 \rangle$$


Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 3, 7 \rangle \cdot \langle 1, -4 \rangle}{\|\langle 3, 7 \rangle\| \|\langle 1, -4 \rangle\|} \quad (\text{given}) \\ &\approx \frac{-25}{(7.616)(4.123)} \approx \frac{-25}{31.401} \quad (\text{by Calc}) \end{aligned}$$

....thus.... $\cos \theta = -0.8$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$)....

$$\begin{aligned} \theta &\approx \cos^{-1}(-0.8) \\ &\approx 2.5 \text{ radians} \\ &\approx 143.2^\circ \end{aligned}$$



(f) Use the dot product to find the 'angle' between the indicated vectors.


$$\vec{w} = \langle 3, 9 \rangle \quad \vec{v} = \langle 7, 1 \rangle$$

Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 3, 9 \rangle \cdot \langle 7, 1 \rangle}{\|\langle 3, 9 \rangle\| \|\langle 7, 1 \rangle\|} \quad (\text{given}) \\ &\approx \frac{30}{(9.487)(7.071)} \approx \frac{30}{67.083} \quad (\text{by Calc}) \end{aligned}$$

$$\dots \text{thus} \dots \quad \cos \theta = 0.45 \quad (\text{by Calc})$$

$$\begin{aligned} \dots \text{then} \dots (\text{assume the sought angle is } 0 \leq \theta \leq 180^\circ) \dots & \quad \theta \approx \cos^{-1}(0.45) \\ & \quad \theta \approx 1.1 \text{ radians} \\ & \quad \approx 63^\circ \end{aligned}$$



(g) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle -5, 2 \rangle \quad \vec{v} = \langle 2, 5 \rangle$$


Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle -5, 2 \rangle \cdot \langle 2, 5 \rangle}{\|\langle -5, 2 \rangle\| \|\langle 2, 5 \rangle\|} \quad (\text{given}) \\ &\approx \frac{0}{(5.385)(5.385)} \approx \frac{0}{28.998} \quad (\text{by Calc}) \end{aligned}$$

....thus.... $\cos \theta = 0$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$)....

$$\begin{aligned} \theta &\approx \cos^{-1}(0) \\ \theta &\approx 1.57 \text{ radians} \\ &\approx 90^\circ \end{aligned}$$



(h) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 1, 7 \rangle \quad \vec{v} = \langle 7, -1 \rangle$$


Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 1, 7 \rangle \cdot \langle 7, -1 \rangle}{\|\langle 1, 7 \rangle\| \|\langle 7, -1 \rangle\|} \quad (\text{given}) \\ &\approx \frac{0}{(7.071)(7.071)} \approx \frac{0}{49.999} \quad (\text{by Calc}) \end{aligned}$$

....thus.... $\cos \theta = 0$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$)....

$$\begin{aligned} \theta &\approx \cos^{-1}(0) \\ \theta &\approx 1.57 \text{ radians} \\ &\approx 90^\circ \end{aligned}$$



- (i) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 2, 3 \rangle \quad \vec{v} = \langle 3, -2 \rangle$$


Solution: Let us assume the angle between the vectors is between 0 and 180° . Suppose we call such angle ' θ ', then....

$$\begin{aligned} \cos \theta &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{v}\| \|\vec{w}\|} \quad (\text{famous DOT product property}) \\ &= \frac{\langle 2, 3 \rangle \cdot \langle 3, -2 \rangle}{\|\langle 2, 3 \rangle\| \|\langle 3, -2 \rangle\|} \quad (\text{given}) \\ &\approx \frac{0}{(3.606)(3.606)} \approx \frac{0}{13.003} \quad (\text{by Calc}) \end{aligned}$$

....thus.... $\cos \theta = 0$ (by Calc)

....then.. (assume the sought angle is $0 \leq \theta \leq 180^\circ$)....

$$\begin{aligned} \theta &\approx \cos^{-1}(0) \\ \theta &\approx 1.57 \text{ radians} \\ &\approx 90^\circ \end{aligned}$$



(j) Use the dot product to find the 'angle' between the indicated vectors.

$$\vec{w} = \langle 1, -3, 5, 0, 6 \rangle \quad \vec{v} = \langle 5, 2, -10, 3, 1 \rangle$$

Solution: don't be afraid, don't google it, don't ask anyone.. just you and the problem.. its on..., if it knocks you down, just get up... don't let it beat you..

6. **Perpendicular Test by the DOT** Find the Dot product for each pair of vectors, then determine if they are perpendicular.

(a) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 1, 2 \rangle \quad \vec{v} = \langle 3, -5 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 3, -5 \rangle \cdot \langle 1, 2 \rangle && \text{(given)} \\ &= -7 && \text{(by inspection)} \end{aligned}$$

...therefore.... \vec{v} and \vec{w} are not perpendicular.

(b) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 3, 2 \rangle \quad \vec{v} = \langle 2, -3 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 2, -3 \rangle \cdot \langle 3, 2 \rangle && \text{(given)} \\ &= 0 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are perpendicular.

- (c) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 1, 4 \rangle \quad \vec{v} = \langle -3, -5 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle -3, -5 \rangle \cdot \langle 1, 4 \rangle && \text{(given)} \\ &= -23 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are not perpendicular.

- (d) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 2, 3 \rangle \quad \vec{v} = \langle 3, -3 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 3, -3 \rangle \cdot \langle 2, 3 \rangle && \text{(given)} \\ &= -3 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are not perpendicular.

- (e) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 3, 6 \rangle \quad \vec{v} = \langle 2, -1 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 2, -1 \rangle \cdot \langle 3, 6 \rangle && \text{(given)} \\ &= 0 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are perpendicular.

- (f) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 3, 12 \rangle \quad \vec{v} = \langle -4, 1 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle -4, 1 \rangle \cdot \langle 3, 12 \rangle && \text{(given)} \\ &= 0 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are perpendicular.

- (g) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle -5, 2 \rangle \quad \vec{v} = \langle 3, 5 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 3, 5 \rangle \cdot \langle -5, 2 \rangle && \text{(given)} \\ &= -5 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are not perpendicular.

- (h) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 1, 7 \rangle \quad \vec{v} = \langle 7, -2 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 7, -2 \rangle \cdot \langle 1, 7 \rangle && \text{(given)} \\ &= -7 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are not perpendicular.

- (i) Use the dot product to find test if the indicated vectors are 'perpendicular'.

$$\vec{w} = \langle 2, 3 \rangle \quad \vec{v} = \langle 3, -2 \rangle$$

Solution: The perpendicular test is quite simple and elegant. Vector \vec{v} is perpendicular to \vec{w} if and only if $\vec{v} \cdot \vec{w} = 0$, or more concisely,

$$\vec{v} \perp \vec{w} \iff \vec{v} \cdot \vec{w} = 0$$

Thus we check..

$$\begin{aligned} \vec{v} \cdot \vec{w} &= \langle 3, -2 \rangle \cdot \langle 2, 3 \rangle && \text{(given)} \\ &= 0 && \text{(by inspection)} \end{aligned}$$

....therefore.... \vec{v} and \vec{w} are perpendicular.

- (j) test to see if perpendicular...

$$\vec{w} = \langle 1, -3, 5, -2, 6 \rangle \quad \vec{v} = \langle 5, 0, -1, 3, 1 \rangle$$

Solution: don't be afraid, don't google it, don't ask anyone.. just you and the problem.. its on..., if it knocks you down, just get up... don't let it beat you..


7. Projections by the DOT

- (a) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 1, 2 \rangle \quad \vec{w} = \langle 5, 0 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 1, 2 \rangle \cdot \langle 5, 0 \rangle}{\|\langle 5, 0 \rangle\|^2} \langle 5, 0 \rangle && \text{(given)} \\ &\approx \frac{5}{25} \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx 0.2 \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx \langle 1, 0 \rangle && \text{(by Calc)} \end{aligned}$$




(b) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 3, 2 \rangle \quad \vec{w} = \langle 5, 0 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 3, 2 \rangle \cdot \langle 5, 0 \rangle}{\|\langle 5, 0 \rangle\|^2} \langle 5, 0 \rangle && \text{(given)} \\ &\approx \frac{15}{25} \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx 0.6 \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx \langle 3, 0 \rangle && \text{(by Calc)} \end{aligned}$$




- (c) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 4, 2 \rangle \quad \vec{w} = \langle 5, 0 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 4, 2 \rangle \cdot \langle 5, 0 \rangle}{\|\langle 5, 0 \rangle\|^2} \langle 5, 0 \rangle && \text{(given)} \\ &\approx \frac{20}{25} \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx 0.8 \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx \langle 4, 0 \rangle && \text{(by Calc)} \end{aligned}$$




- (d) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 7, 2 \rangle \quad \vec{w} = \langle 5, 0 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 7, 2 \rangle \cdot \langle 5, 0 \rangle}{\|\langle 5, 0 \rangle\|^2} \langle 5, 0 \rangle && \text{(given)} \\ &\approx \frac{35}{25} \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx 1.4 \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx \langle 7, 0 \rangle && \text{(by Calc)} \end{aligned}$$




(e) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle -3, 2 \rangle \quad \vec{w} = \langle 5, 0 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle -3, 2 \rangle \cdot \langle 5, 0 \rangle}{\|\langle 5, 0 \rangle\|^2} \langle 5, 0 \rangle && \text{(given)} \\ &\approx \frac{-15}{25} \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx -0.6 \langle 5, 0 \rangle && \text{(by Calc)} \\ &\approx \langle -3, 0 \rangle && \text{(by Calc)} \end{aligned}$$




(f) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 1, 2 \rangle \quad \vec{w} = \langle 0, 6 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 1, 2 \rangle \cdot \langle 0, 6 \rangle}{\|\langle 0, 6 \rangle\|^2} \langle 0, 6 \rangle && \text{(given)} \\ &\approx \frac{12}{36} \langle 0, 6 \rangle && \text{(by Calc)} \\ &\approx 0.333 \langle 0, 6 \rangle && \text{(by Calc)} \\ &\approx \langle 0, 2 \rangle && \text{(by Calc)} \end{aligned}$$




(g) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 3, 2 \rangle \quad \vec{w} = \langle 0, 7 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 3, 2 \rangle \cdot \langle 0, 7 \rangle}{\|\langle 0, 7 \rangle\|^2} \langle 0, 7 \rangle && \text{(given)} \\ &\approx \frac{14}{49} \langle 0, 7 \rangle && \text{(by Calc)} \\ &\approx 0.286 \langle 0, 7 \rangle && \text{(by Calc)} \\ &\approx \langle 0, 2 \rangle && \text{(by Calc)} \end{aligned}$$




(h) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 4, 2 \rangle \quad \vec{w} = \langle 0, 2 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 4, 2 \rangle \cdot \langle 0, 2 \rangle}{\|\langle 0, 2 \rangle\|^2} \langle 0, 2 \rangle && \text{(given)} \\ &\approx \frac{4}{4} \langle 0, 2 \rangle && \text{(by Calc)} \\ &\approx 1 \langle 0, 2 \rangle && \text{(by Calc)} \\ &\approx \langle 0, 2 \rangle && \text{(by Calc)} \end{aligned}$$




- (i) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 7, 2 \rangle \quad \vec{w} = \langle 0, 4 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 7, 2 \rangle \cdot \langle 0, 4 \rangle}{\|\langle 0, 4 \rangle\|^2} \langle 0, 4 \rangle && \text{(given)} \\ &\approx \frac{8}{16} \langle 0, 4 \rangle && \text{(by Calc)} \\ &\approx 0.5 \langle 0, 4 \rangle && \text{(by Calc)} \\ &\approx \langle 0, 2 \rangle && \text{(by Calc)} \end{aligned}$$




- (j) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle -3, 2 \rangle \quad \vec{w} = \langle 0, 3 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle -3, 2 \rangle \cdot \langle 0, 3 \rangle}{\|\langle 0, 3 \rangle\|^2} \langle 0, 3 \rangle && \text{(given)} \\ &\approx \frac{6}{9} \langle 0, 3 \rangle && \text{(by Calc)} \\ &\approx 0.667 \langle 0, 3 \rangle && \text{(by Calc)} \\ &\approx \langle 0, 2 \rangle && \text{(by Calc)} \end{aligned}$$




(k) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 1, 2 \rangle \quad \vec{w} = \langle 1, 6 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 1, 2 \rangle \cdot \langle 1, 6 \rangle}{\|\langle 1, 6 \rangle\|^2} \langle 1, 6 \rangle && \text{(given)} \\ &\approx \frac{13}{37} \langle 1, 6 \rangle && \text{(by Calc)} \\ &\approx 0.351 \langle 1, 6 \rangle && \text{(by Calc)} \\ &\approx \langle 0.35, 2.11 \rangle && \text{(by Calc)} \end{aligned}$$




- (l) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 3, 2 \rangle \quad \vec{w} = \langle 7, 1 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 3, 2 \rangle \cdot \langle 7, 1 \rangle}{\|\langle 7, 1 \rangle\|^2} \langle 7, 1 \rangle && \text{(given)} \\ &\approx \frac{23}{50} \langle 7, 1 \rangle && \text{(by Calc)} \\ &\approx 0.46 \langle 7, 1 \rangle && \text{(by Calc)} \\ &\approx \langle 3.22, 0.46 \rangle && \text{(by Calc)} \end{aligned}$$




- (m) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 4, 2 \rangle \quad \vec{w} = \langle 4, 2 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 4, 2 \rangle \cdot \langle 4, 2 \rangle}{\|\langle 4, 2 \rangle\|^2} \langle 4, 2 \rangle && \text{(given)} \\ &\approx \frac{20}{20} \langle 4, 2 \rangle && \text{(by Calc)} \\ &\approx 1 \langle 4, 2 \rangle && \text{(by Calc)} \\ &\approx \langle 4, 2 \rangle && \text{(by Calc)} \end{aligned}$$




(n) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 7, 2 \rangle \quad \vec{w} = \langle 8, 1 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 7, 2 \rangle \cdot \langle 8, 1 \rangle}{\|\langle 8, 1 \rangle\|^2} \langle 8, 1 \rangle && \text{(given)} \\ &\approx \frac{58}{65} \langle 8, 1 \rangle && \text{(by Calc)} \\ &\approx 0.892 \langle 8, 1 \rangle && \text{(by Calc)} \\ &\approx \langle 7.14, 0.89 \rangle && \text{(by Calc)} \end{aligned}$$




- (o) Use the dot product to find the 'projection' of \vec{v} onto \vec{w} then draw such projection.

$$\vec{v} = \langle 3, 2 \rangle \quad \vec{w} = \langle 9, 3 \rangle$$

Solution:

$$\begin{aligned} \text{proj}_{\vec{w}} \vec{v} &= \frac{\vec{v} \cdot \vec{w}}{\|\vec{w}\|^2} \vec{w} && \text{(famous DOT product property)} \\ &= \frac{\langle 3, 2 \rangle \cdot \langle 9, 3 \rangle}{\|\langle 9, 3 \rangle\|^2} \langle 9, 3 \rangle && \text{(given)} \\ &\approx \frac{33}{90} \langle 9, 3 \rangle && \text{(by Calc)} \\ &\approx 0.367 \langle 9, 3 \rangle && \text{(by Calc)} \\ &\approx \langle 3.3, 1.1 \rangle && \text{(by Calc)} \end{aligned}$$



- (p) find projection

$$\vec{v} = \langle 1, -3, 5, -2, 6 \rangle \quad \text{onto } \vec{w} = \langle 5, 0, -1, 3, 1 \rangle$$

Solution: don't be afraid, don't google it, don't ask anyone.. just you and the problem.. its on..., if it knocks you down, just get up... don't let it beat you..